

Analytic decomposition of differential graded Lie algebras

Frank Schuhmacher

February 2, 2008

Abstract

We prove explicit formulas for the decomposition of a differential graded Lie algebra into a minimal and a linear L_∞ -algebra. We define a category of metric L_∞ -algebras, called Palamodov L_∞ algebras, where the structure maps satisfy a certain convergence condition and deduce a decomposition theorem for differential graded Lie algebras in this category. This theorem serves for instance to prove the convergence of the Kuranishi map assigned to a differential graded Lie algebra.

Contents

1	Formal DG manifolds	3
1.1	Graded symmetric and exterior algebras	3
1.2	Free differential graded coalgebras	4
1.3	Formal vector fields as coderivations	6
1.4	Formal DG manifolds and L_∞ -algebras	8
2	Trees	9
2.1	Definitions	10
2.2	Operations on trees	12
3	Decomposition of differential graded Lie algebras	13
4	Palamodov DG manifolds	20
4.1	Palamodov spaces	20
4.2	Decomposition of Palamodov DG Lie algebras	24

Introduction

As observed by Kontsevich [8] and others, any L_∞ -algebra over a field of characteristic zero can be decomposed into a direct sum of a minimal and a linear contractible L_∞ -algebra. This means that there exists an L_∞ -structure on the cohomology $H(L)$

of the complex (L, d) , where d is the linear component of the L_∞ -structure of L , such that $H(L)$ is a direct summand of L , not only as a graded module but also as an L_∞ -algebra. This has two important consequences: first, any quasi-isomorphism between L_∞ -algebras over a field of characteristic zero can be inverted (we may thus speak of L_∞ -equivalence) and secondly, the deformation functors (local Artin rings) $\rightarrow$ (sets) assigned to quasi-isomorphic DG Lie algebras (see [8] or [13]) are isomorphic. Thus, if the deformations of an object are *governed* by a DG Lie algebra L (i.e. deformations of the object correspond to Maurer-Cartan elements of L and equivalent deformations are in the same orbit of the action of a Lie group related to L^0), then the L_∞ -equivalence class of L contains the formal deformation theory of the object. For example, the base of a formally versal deformation, if it exists, is defined by the *Kuranishi-map* $H(L)^1 \rightarrow H(L)^2$, whose Taylor coefficients are the restrictions of the L_∞ -structure maps of $H(L)$. This motivates the question of an explicit description of the L_∞ -structure on $H(L)$ and of an L_∞ -map $H(L) \rightarrow L$ in the case where L is just a DG Lie algebra (recursion formulas were already indicated by Kontsevich [7] and Huebschmann/Stasheff [4]). We give an explicit description for the structure on $H(L)$ in function of a splitting (i.e. a degree -1 map $\eta : L \rightarrow L$ with $d\eta d = d$) of the complex (L, d) , and an explicit description of an L_∞ -isomorphism $H(L) \oplus (F, d) \rightarrow L$, where F is the complement of $H(L)$ in L . The improvement with respect to [20] is that now, we also have an explicit description of the map $(F, d) \rightarrow L$. In [20], the existence of such a map was obtained by obstruction theory. Since an inverse isomorphism can recursively be constructed by the inverse function theorem for L_∞ -algebras, this allows a complete algorithmic decomposition of a split DG Lie algebra.

There is the deeper question of the convergence of the Kuranishi map. The present paper contains a quite general answer to this question: To formulate the main result, we need some analytic language: First of all, the algebraic language of L_∞ -algebras or equivalently, of formal DG manifolds, is not sufficient for convergence questions. It would be straight forward, but for many deformation problems still not sufficient, to define Banach L_∞ -algebras, i.e. L_∞ -structures $(\mu_n)_{n \in \mathbb{N}}$ on underlying graded Banach spaces L , such that the formal sums $\sum_n \frac{1}{n!} \tilde{\mu}_n$, with $\tilde{\mu}_n(x) = \mu_n(x, \dots, x)$ converge on some zero neighbourhood. We introduce, more generally, *Palamodov L_∞ -algebras*, where the underlying graded vector spaces do not even have a norm but a metric defined by a countable family of semi-norms. We need a suitable notion of convergence, for this context. For *Palamodov DG Lie algebras* admitting a splitting, we prove the convergence of the structure μ on $H(L)$ and of the map $H(L) \oplus F \rightarrow L$ and since the inverse function theorem is valid for Palamodov L_∞ -algebras, this implies the decomposition

$$(L, d, [\cdot, \cdot]) \cong (H, \mu) \oplus (F, d)$$

in the category of Palamodov L_∞ -algebras. Recall that this decomposition specialises to the Banach context.

1 Formal DG manifolds

In this section, we give a detailed review of the correspondence between L_∞ -algebras, formal DG-manifolds and free DG coalgebras. Experts may skip this section.

1.1 Graded symmetric and exterior algebras

Let $\mathbb{K}$ be a commutative ring of characteristic zero. For a graded $\mathbb{K}$ -module M , the graded symmetric algebra SM is defined as the tensor algebra $TM = \bigoplus_{n \geq 0} M^{\otimes n}$ modulo the relations $m_1 \otimes m_2 - (-1)^{m_1 m_2} m_2 \otimes m_1 = 0$. We denote the graded symmetrical product by $\odot$. The algebra TM (resp. SM) is bigraded. The graduation on TM (resp. SM) defined by $g(m_1 \otimes \dots \otimes m_n) = g(m_1) + \dots + g(m_n)$ (resp. $g(m_1 \odot \dots \odot m_n) = g(m_1) + \dots + g(m_n)$), where g is the graduation of M , will be called **linear graduation**. The one defined by $g(m_1 \otimes \dots \otimes m_n) = n$ (resp. $g(m_1 \odot \dots \odot m_n) = n$) will be called **polynomial graduation**. On $S_+M := \bigoplus_{n \geq 1} M^{\odot n}$, there is a natural $\mathbb{K}$ -linear comultiplication $\Delta^+ : S_+M \rightarrow S_+M \otimes S_+M$, given by

$$m_1 \odot \dots \odot m_n \mapsto \sum_{j=1}^{n-1} \sum_{\sigma \in \text{Sh}(j, n)} \epsilon(\sigma, m_1, \dots, m_n) m_{\sigma(1)} \odot \dots \odot m_{\sigma(j)} \otimes m_{\sigma(j+1)} \odot \dots \odot m_{\sigma(n)},$$

where the sign $\epsilon(\sigma) := \epsilon(\sigma, m_1, \dots, m_n)$ is defined in such a way that $m_{\sigma(1)} \odot \dots \odot m_{\sigma(n)} = \epsilon(\sigma) m_1 \odot \dots \odot m_n$. Note that $\text{Kern } \Delta^+ = M$. On SM , there is a $\mathbb{K}$ -linear comultiplication Δ , defined by $\Delta(1) := 1 \otimes 1$ and $\Delta(m) := m \otimes 1 + \Delta'(w) + 1 \otimes m$, for $m \in S_+M$. Note that Δ is injective.

For a graded module L , the graded exterior algebra $\bigwedge^+ L$ without unit is defined as the tensor algebra $T_+L = \bigoplus_{n \geq 1} L^{\otimes n}$ modulo the relations $a_1 \otimes a_2 + (-1)^{a_1 a_2} a_2 \otimes a_1 = 0$. We denote the graded exterior product by $\wedge$. By $L[1]$, we denote the graded module with $L[1]^i = L^{i+1}$ and by $\downarrow$ the canonical map $L \rightarrow L[1]$ of degree -1 . Set $\uparrow := \downarrow^{-1}$. Remark that for each $n \geq 1$, there is an isomorphism

$$\begin{aligned} \downarrow^n : \bigwedge^n L &\longrightarrow L[1]^{\odot n} \\ a_1 \wedge \dots \wedge a_n &\mapsto (-1)^{(n-1) \cdot a_1 + \dots + 1 \cdot a_{n-1}} \downarrow a_1 \odot \dots \odot \downarrow a_n. \end{aligned}$$

Its inverse map is given by $(-1)^{\frac{n(n-1)}{2}} \uparrow^n$. In this formula, we deduce the sign from the Koszul convention. More generally, for homogeneous graded morphisms f, g of graded modules, we set $(f \otimes g)(a \otimes b) := (-1)^{g a} f(a) \otimes g(b)$. In the exponent, a always means the degree of an homogeneous element (or morphism) a and ab means the product of degrees and not the degree of the product.

For $\sigma \in \Sigma_n$ and $a_1, \dots, a_n \in L$, we define the sign $\chi(\sigma) := \chi(\sigma, a_n, \dots, a_1)$ in such a way that

$$a_{\sigma(1)} \wedge \dots \wedge a_{\sigma(n)} = \chi(\sigma) a_1 \wedge \dots \wedge a_n.$$

The correlation between χ and ϵ is given by

$$\chi(\sigma, a_1, \dots, a_n) = (-1)^{(n-1)(a_1 + a_{\sigma(1)}) + \dots + 1 \cdot (a_{n-1} + a_{\sigma(n-1)})} \epsilon(\sigma, \downarrow a_1, \dots, \downarrow a_n),$$

for $a_1, \dots, a_n \in L$. For a graded module V , we define two different actions of the symmetric group Σ_n on $V^{\otimes n}$: The first one is given by

$$(\sigma, v_1 \otimes \dots \otimes v_n) \mapsto \epsilon(\sigma, v_1, \dots, v_n) v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(n)}.$$

The application of a σ commutes with the canonical projection $V^{\otimes n} \rightarrow V^{\odot n}$. The second one is given by

$$(\sigma, v_1 \otimes \dots \otimes v_n) \mapsto \chi(\sigma, v_1, \dots, v_n) v_{\sigma(1)} \otimes \dots \otimes v_{\sigma(n)}.$$

Here, the application of a σ commutes with the canonical projection $V^{\otimes n} \rightarrow \wedge^n V$. When we work with symmetric powers, we use the first action; when we work with exterior powers, we use the second one. Since the context shall always be clear, we don't distinguish both actions by different notation. We will use the (anti-)symmetrization maps:

$$\alpha_n := \sum_{\sigma \in \Sigma_n} \sigma. : V^{\otimes n} \longrightarrow V^{\otimes n}.$$

When $\sigma.$ denotes the first action, α_n can be considered as a map $V^{\odot n} \rightarrow V^{\otimes n}$; when $\sigma.$ denotes the second action, α_n can be considered as a map $\wedge^n V \rightarrow V^{\otimes n}$. Furthermore, for both cases, we will use the maps

$$\alpha_{k,n} := \sum_{\sigma \in \text{Sh}(k,n)} \sigma. : V^{\otimes n} \longrightarrow V^{\otimes n}.$$

For the natural projection $\pi : V^{\otimes n} \rightarrow V^{\odot n}$ (resp. $V^{\otimes n} \rightarrow \wedge^n V$), we have

$$\pi \circ \alpha = n! \text{Id}.$$

1.2 Free differential graded coalgebras

Let (C_1, Δ_1) and (C_2, Δ_2) be coalgebras. Remember that a module homomorphism $F : C_1 \rightarrow C_2$ is a coalgebra morphism, if and only if the diagram

$$\begin{array}{ccc} C_1 & \xrightarrow{F} & C_2 \\ \downarrow \Delta_1 & & \downarrow \Delta_2 \\ C_1 \otimes C_1 & \xrightarrow{F \otimes F} & C_2 \otimes C_2 \end{array} \quad (1.1)$$

commutes. Each coalgebra morphism $F : (SM, \Delta) \rightarrow (SM', \Delta)$ satisfies $F(1) = 1$. The restriction $F \mapsto F|_{S_+M}$ is a one-to-one correspondence between coalgebra morphisms $(SM, \Delta) \rightarrow (SM', \Delta)$ and coalgebra morphisms $F : (S_+M, \Delta^+) \rightarrow (S_+M', \Delta^+)$. We fix the following notions:

$$\begin{aligned} \hat{F}_n &:= F|_{M^{\odot n}} : M^{\odot n} \longrightarrow SM' \\ F_{k,l} &:= \text{pr}_{M'^{\odot l}} \circ \hat{F}_k : M^{\odot k} \longrightarrow M'^{\odot l} \\ F_n &:= F_{n,1} : M^{\odot n} \longrightarrow M' \end{aligned}$$

Sometimes, we shall consider the maps F_n as graded symmetric maps $M^{\otimes n} \rightarrow M'$ instead of maps $M^{\odot n} \rightarrow M'$. For each multi-index $I = (i_1, \dots, i_k) \in \mathbb{N}^k$, we set $I! := i_1! \cdot \dots \cdot i_k!$ and $|I| := i_1 + \dots + i_k$ and

$$F_I := \frac{1}{I!k!} (F_{i_1} \odot \dots \odot F_{i_k}) \circ \alpha_n.$$

Here, by $F_{i_1} \odot \dots \odot F_{i_k}$, we mean the composition of $F_{i_1} \otimes \dots \otimes F_{i_k}$ and the natural projection $M'^{\otimes k} \rightarrow M'^{\odot k}$. A morphism $F : SM \rightarrow SM$ of graded coalgebras is called **strict**, if $F_n = 0$ for each $n \geq 2$.

For a coalgebra (C, Δ) , remember that a module homomorphism $Q : C \rightarrow C$ is a coderivation, if and only if the diagram

$$\begin{array}{ccc} C & \xrightarrow{Q} & C \\ \downarrow \Delta & & \downarrow \Delta \\ C \otimes C & \xrightarrow{Q \otimes 1 + 1 \otimes Q} & C \otimes C \end{array} \quad (1.2)$$

commutes. We fix the following notions:

$$\begin{aligned} \hat{Q}_n &:= Q|_{M^{\odot n}} : M^{\odot n} \longrightarrow SM \\ Q_{k,l} &:= \text{pr}_{M^{\odot l}} \circ \hat{Q}_k : M^{\odot k} \longrightarrow M^{\odot l} \\ Q_n &:= Q_{n,1} : M^{\odot n} \longrightarrow M \end{aligned}$$

Observe that there is a one-to-one correspondence between degree i coderivations on (S_+M, Δ^+) and degree i coderivations of Q degree $+1$ on (SM, Δ) with $Q(1) = 0$. By the following proposition, there is a one-to-one correspondence between degree i coderivations $Q : SM \rightarrow SM$ and sequences of linear maps $Q_n : S_nM \rightarrow M$ of degree i , and a one-to-one correspondence between coalgebra maps $F : SM \rightarrow SM'$ and sequences of linear maps $F_n : S_nM \rightarrow M'$, $n \geq 1$.

Proposition 1.1. *Let Q (resp. Q') be a degree i coderivation on SM (resp. SM') and $F : SM \rightarrow SM'$ a morphism of graded coalgebras. Then, for $n \geq 1$ and $1 \leq l \leq n+1$, $1 \leq k \leq n$, we have*

$$Q_{n,l} = (Q_{n-l+1} \otimes 1 \otimes \dots \otimes 1) \circ \alpha_{n-l+1,n}$$

and

$$F_{n,k} = \sum_{i_1 + \dots + i_k = n} F_I.$$

The map F respects the coderivations Q and Q' if and only if $F(Q(1)) = Q'(1)$ and for each $n \geq 1$, we have

$$\sum_{k=1}^n \sum_{\substack{I \in \mathbb{N}^k \\ |I|=n}} Q'_k \circ F_I = \sum_{k+l=n+1} F_l \circ (Q_k \otimes 1 \otimes \dots \otimes 1) \circ \alpha_{k,n}. \quad (1.3)$$

On the right hand-side, the sum is over all $l \geq 1$ and $k \geq 0$. The term $(Q_0 \otimes 1 \otimes \dots \otimes 1)(m_1 \otimes \dots \otimes m_n)$ must be interpreted as $Q_0(1) \otimes m_1 \otimes \dots \otimes m_n$.

Set $\text{Coder}^i(SM)$ to be the module of degree i coderivations. The direct sum $\text{Coder}(SM) = \bigoplus_{i \in \mathbb{Z}} \text{Coder}^i(SM)$ is a graded Lie algebra with graded commutator

$$[Q, q] = Q \circ q - (-1)^{Q \cdot q} q \circ Q$$

as Lie bracket. Explicitly,

$$[Q, q]_n = \sum_{k+l=n+1} Q_l(q_k \otimes 1^{\otimes n-k}) \circ \alpha_{k,n} - (-1)^{Q \cdot q} q_l \circ (Q_k \otimes 1^{\otimes n-k}) \circ \alpha_{k,n}. \quad (1.4)$$

Observe that a degree 1 coderivation Q on SM is a codifferential (i.e. $Q^2 = 0$) if and only if, for each $n \geq 0$, we have

$$\sum_{k+l=n+1} Q_l(Q_k \otimes 1^{\otimes n-k}) \circ \alpha_{k,n} = 0.$$

By definition, a free DG coalgebra is a pair (SM, Q^M) , where M is a graded $\mathbb{K}$ -module and Q^M is a degree one codifferential on SM . A morphism of free DG coalgebras is a coalgebra map of degree zero respecting the codifferentials.

1.3 Formal vector fields as coderivations

Before we introduce the equivalent notions of formal DG manifolds and L_∞ -algebras, we make some remarks on analytic space germs and on analytic vector fields: An analytic function $f : \mathbb{C}^N \longrightarrow \mathbb{C}^m$ defined on a neighbourhood of 0 can be represented by a power series

$$f(x) = \sum_{\nu} a_{\nu} x^{\nu}$$

with coefficients a_{ν} in $\mathbb{C}^m$ or as a sum

$$f(x) = \sum_{n=0}^{\infty} \frac{1}{n!} \tilde{f}_n(x),$$

where $\tilde{f}_n(x) = f_n(x, \dots, x)$, for a uniquely defined symmetric n -linear map $f_n : \mathbb{C}^N \otimes \dots \otimes \mathbb{C}^N \longrightarrow \mathbb{C}^m$. The map $\tilde{f}_n$ is the n -th Taylor coefficient of f . The relation between the coefficients a_{ν} and f_n is given by the formulas

$$f_n = \sum_{|\nu|=n} a_{\nu} (e^*)^{\odot \nu}$$

and

$$a_{\nu} = \frac{1}{n!} f_{|\nu|}(e^{\odot \nu}).$$

For the verification of this formulas, remark that

$$(a_\nu(e^*)^{\odot \nu})^\sim(x) = \frac{|\nu|!}{\nu!} a_\nu(e^*)^{\odot \nu}(x^{\odot \nu}) = |\nu|! a_\nu x^\nu.$$

Let $R = \mathbb{C}\langle e_j \rangle_{j \in J}$ be a free g-finitely generated algebra over the convergent power series algebra $R^0 = \mathbb{C}\{e_j\}_{j \in J^0}$. Let $\hat{R} = \mathbb{C}[[x_j]]$ be the $\mathfrak{m}$ -adic completion of R , where $\mathfrak{m}$ is the ideal generated by the x_j ; $j \in J$. Consider the graded $\mathbb{C}$ -vector space $M = \coprod_{j \in J} \mathbb{C}e_j$ with generators of degree $-g(e_j) = g(x_j)$. Consider SM as graded coalgebra. The multiplication $\odot$ in the dual algebra $(SM)^*$ of (SM, Δ) is given by

$$(a \odot b)(m) = (a \otimes b)(\Delta(m)) = \sum_{\sigma \in \text{Sh}(k, k+l)} \epsilon(\sigma, m) (-1)^{m'_\sigma \cdot b} a(m'_\sigma) \cdot b(m''_\sigma),$$

for $a \in (M^{\odot k})^*$, $b \in (M^{\odot l})^*$ and $m \in M^{\odot k+l}$. We have an isomorphism $\hat{R} \rightarrow (SM)^*$ of graded algebras sending x_j to the dual generator e_j^* of M^* . The inverse map is given by

$$f \mapsto \sum_{n=0}^{\infty} \sum_{|\nu|=n} \frac{1}{\nu!} f_n(e^{\odot \nu}) \cdot x^\nu,$$

where we sum only over the ordered multi-indices with indices of odd degree appearing at most once. For an ordered multi-index $\nu = (\nu_1, \dots, \nu_N)$, we have

$$(e^*)^{\odot \nu} := (e_1^*)^{\odot \nu_1} \odot \dots \odot (e_N^*)^{\odot \nu_N} = \nu! (e^{\odot \nu})^*.$$

Let $\mathfrak{m}$ be the canonical maximal ideal of $\hat{R}$ and set $\text{Der}(\hat{R}) = \oplus_{i \in \mathbb{Z}} \text{Der}^i(\hat{R})$ to be the graded vector space of all $\mathbb{C}$ -linear, $\mathfrak{m}$ -adic continuous derivations of $\hat{R}$. The graded commutator defines a graded Lie structure on $\text{Der}(\hat{R})$. Observe that the dualization map $Q \mapsto Q^*$ is an isomorphism of graded Lie algebras

$$\text{Coder}(SM) \longrightarrow \text{Der}(S(M)^*).$$

The inverse map sends a derivation δ to the sum $\sum_{j \in J} e_j \cdot \delta(e_j^*)$.

Consider an open subset M of $\mathbb{C}^m$ with $0 \in M$. An analytic vectorfield q on M is an analytic map $M \rightarrow \mathbb{C}^m$. Locally at 0, it is given by a power series

$$\tilde{q} : x \mapsto \sum_{n=0}^{\infty} \frac{1}{n!} q_n(x, \dots, x),$$

where $q_n : \mathbb{C}^m \times \dots \times \mathbb{C}^m \rightarrow \mathbb{C}^m$ is a unique symmetric n-linear map, i.e. $\tilde{q}_n : x \mapsto q_n(x, \dots, x)$ is a homogeneous polynomial function of degree n . Recall that as a Lie algebra, we can identify the set of analytic vectorfields on M with the Lie algebra of $\mathbb{C}$ -linear derivations of the section algebra $\Gamma(\mathcal{O}_M, M)$, where the Lie bracket is the commutator of derivations. The following exercise gives the algebraic description of this Lie bracket:

Exercise 1. For vectorfields Q, q on M , we have $[Q, q](x) = \sum_{n=0}^{\infty} \frac{1}{n!} [Q, q]_n^{\sim}(x)$, where the symmetric n -linear map $[Q, q]_n$ is given by the following formula:

$$[Q, q]_n(x_1, \dots, x_n) = \sum_{k+l=n+1} \sum_{\sigma \in \text{Sh}(k, n)} Q_l(q_k(x_{\sigma(1)}, \dots, x_{\sigma(k)}), x_{\sigma(k+1)}, \dots, x_{\sigma(n)}) - q_l(Q_k(x_{\sigma(1)}, \dots, x_{\sigma(k)}), x_{\sigma(k+1)}, \dots, x_{\sigma(n)}). \quad (1.5)$$

If we replace M by an arbitrary vector space, we can define the Lie algebra of **formal vectorfields** on M , i.e. of formal sums $q = \sum \frac{1}{n!} \tilde{q}_n$, where each $q_n : M^{\otimes n} \rightarrow M$ is a symmetric n -linear map and the bracket $[Q, q]$ of formal vectorfields Q, q on M is the formal sum $\sum [Q, q]_n$, where $[Q, q]_n$ is defined by the equation (1.5). To define formal DG manifolds, we generalise the definition of formal vectorfields to the graded context, where the word *symmetry* is replaced by the word *graded symmetry*.

1.4 Formal DG manifolds and L_{∞} -algebras

Let M be a graded $\mathbb{K}$ -module. A **formal vectorfield of degree $i \in \mathbb{Z}$** on M is a formal sum $q = \sum \frac{1}{n!} \tilde{q}_n$, where each $q_n : M^{\otimes n} \rightarrow M$ is a graded symmetric n -linear map of linear degree i . Set $\text{Vect}^i(M)$ to be the $\mathbb{K}$ -module of formal vectorfields of degree i on M . The direct sum $\text{Vect}(M) := \sum_{i \in \mathbb{Z}} \text{Vect}^i(M)$ has a graded Lie algebra structure with Lie bracket defined by equation (1.4). By definition, the graded Lie algebras $\text{Vect}(M)$ and $\text{Coder}(SM)$ are canonically isomorphic. A **formal DG manifold** is a pair (M, Q^M) , where M is a graded $\mathbb{K}$ -module and Q^M is a formal degree one vectorfield on M with $[Q^M, Q^M] = 0$. In other words, a formal DG manifold is a pair (M, Q^M) such that the pair (SM, Q^M) is a free DG coalgebra. A **morphism** $f : M \rightarrow M'$ of formal DG manifolds is a formal sum $f = \sum_{n=0}^{\infty} \tilde{f}_n$, where $f_n : M^{\otimes n} \rightarrow M'$ is a graded symmetric map such that the family $(f_n)_{n \geq 1}$ defines a morphism $SM \rightarrow SM'$ of DG coalgebras. Thus, the category $DG\text{-MANF}$ of formal DG manifolds is canonically isomorphic to the category of free DG coalgebras. Geometrically, a morphism of formal DG manifolds is a morphism of space germs, equivariant with respect to the given vectorfields.

Remember that a module L with a sequence of maps $\mu_n : \bigwedge^n L \rightarrow L$ of degree $2 - n$, for $n \geq 0$, is called an **L_{∞} -algebra** if the pair (M, Q^M) with $M := L[1]$, and

$$Q_n := (-1)^{n(n-1)/2} \downarrow \circ \mu_n \circ \uparrow^n : M^{\odot n} \rightarrow M$$

is a formal DG manifold. This condition is equivalent to the equations

$$\sum_{k+l=n+1} \sum_{\sigma \in \text{Sh}(k, n)} (-1)^{k(l-1)} \chi(\sigma) \mu_l(\mu_k(a_{\sigma(1)}, \dots, a_{\sigma(k)}), a_{\sigma(k+1)}, \dots, a_{\sigma(n)}) = 0 \quad (1.6)$$

for each $n \geq 0$ and $a_1, \dots, a_n \in L$. In the literature, μ_0 is mostly assumed to be trivial. If this is the case, (L, μ_1) is a DG module. An L_{∞} -algebra $(L, \mu_n)_{n \geq 1}$ is called **minimal**, if $\mu_1 = 0$. It is called **linear**, if $\mu_i = 0$ for $i \geq 2$.

Let (L, μ_*) and (L', μ'_*) be L_∞ -algebras. Set $M := L[1]$, $M' := L'[1]$. A sequence of maps $f_n : \bigwedge^n L \longrightarrow L'$; $n \geq 1$ of degree $1 - n$ is called **L_∞ -morphism**, if the maps $F_n := W^{\odot n} \longrightarrow W'$ induced by f_n (explicitly: $F_n = (-1)^{n(n-1)/2} \downarrow \circ f_n \circ \uparrow^n$) define a morphism of formal DG manifolds. Rewrite condition (1.3) in terms of f_n and μ_n :

$$\begin{aligned} & \mu'_1 \circ f_n - \sum_{i+j=n} \frac{(-1)^i}{2} \mu'_2(f_i, f_j) \circ \alpha_{i,j} \\ & + \sum_{k=3}^n \sum_{\substack{I \in \mathbb{N}^k \\ |I|=n}} (-1)^{k(k-1)/2 + i_1(k-1) + \dots + i_{k-1} \cdot 1} \mu'_k \circ f_I \\ & = \sum_{k+l=n+1} (-1)^{k(l-1)} f_l \circ (\mu_k \otimes 1 \otimes \dots \otimes 1) \circ \alpha_{k,n} \end{aligned} \quad (1.7)$$

For the case where L' is a differential graded Lie-algebra, i.e. $\mu'_k = 0$ for $k = 0$ and $k \geq 3$, set $d := \mu'_1$ and $[\ , \] := \mu'_2$. We get the following conditions for the maps f_n to define an L_∞ -morphism (see Definition 5.2 of [11]):

$$\begin{aligned} & df_n(a_1, \dots, a_n) - \\ & \sum_{i+j=n} \sum_{\sigma} \chi(\sigma) (-1)^{i+(j-1)(a_{\sigma(1)} + \dots + a_{\sigma(i)})} [f_i(a_{\sigma(1)}, \dots, a_{\sigma(i)}), f_j(a_{\sigma(i+1)}, \dots, a_{\sigma(n)})] \\ & = \sum_{k+l=n+1} \sum_{\sigma \in \text{Sh}(k,n)} (-1)^{k(l-1)} \chi(\sigma) f_l(\mu_k(a_{\sigma(1)}, \dots, a_{\sigma(k)}), a_{\sigma(k+1)}, \dots, a_{\sigma(n)}), \end{aligned}$$

where $a_1, \dots, a_n \in L$ and the second sum goes over all σ in $\text{Sh}(i, n)$ such that $\sigma(1) < \sigma(i+1)$. A morphism $f : (L, \mu_n)_{n \geq 1} \longrightarrow (L', \mu'_n)_{n \geq 1}$ of L_∞ -algebras is called an **L_∞ -quasi-isomorphism**, if f_1 is a quasi-isomorphism of differential graded modules.

Proposition 1.2. *Let $f : M \longrightarrow M'$ be a morphism of formal supermanifolds. Suppose, there is a module homomorphism $g' : M' \longrightarrow M$ such that $g' \circ f_1 = \text{Id}_M$. Then, there is a morphism $g : M' \longrightarrow M$ of formal supermanifolds such that $g_1 = g'$ and $gf = \text{Id}_M$. If f_1 is an isomorphism with inverse g and if f is Q -equivariant and if g' respects $Q_1^{M'}$ and Q_1^M , then g can be chosen Q -equivariant as well.*

Proof. One can check directly that the sequence of maps defined by

$$g_n := - \sum_{k=2}^n \sum_{\substack{I \in \mathbb{N}^k \\ |I|=n}} g_1 \circ f_k \circ (g_I),$$

for $n \geq 2$, define a morphism of formal supermanifolds with the desired property. $\square$

2 Trees

Trees were first used by Kontsevich/ Soibelman [9] to describe the A_∞ -structure that a DG module, homotopy equivalent to a differential graded algebra inherits. We have a similar objective, but for L_∞ -algebras instead of A_∞ -algebras. Trees in our definition are always binary trees. We give a definition of binary trees and assign several invariants to them, which are important in order to get good signs later on.

2.1 Definitions

A **(binary) tree** with n leaves consists of a pair $\phi = (\phi, V)$ where $V = \{K_0, \dots, K_{n-2}\}$ denotes a set of **ramifications** and $\phi : \{K_1, \dots, K_{n-1}\} \longrightarrow \{K_0, \dots, K_{n-1}\}$ is a map, such that for each $i = 0, \dots, n-2$,

- (i) the inverse image $\phi^{-1}(K_i)$ contains at most 2 elements;
- (ii) there is an $n \geq 0$ such that $\phi^n(K_i) = K_0$.

K_0 is called **root** of ϕ . There is a tree with one leaf and no ramification which will always be denoted by τ . An **orientation** of a tree (ϕ, V) is a family $\pi = (\pi_K)_{K \in V}$ of inclusions $\pi_K : \phi^{-1}(K) \longrightarrow \{1, 2\}$. The triple $\phi = (\phi, V, \pi)$ is called an **oriented tree**. For each oriented tree (ϕ, V, π) , there is a natural **ordering** on the set V : For $K \in V \setminus K_0$, suppose that $\phi^m(K) = K_0$. We set

$$v(K) := \frac{\pi_{\phi(K)}(K)}{3^m} + \frac{\pi_{\phi^2(K)}(\phi(K))}{3^{m-1}} + \dots + \frac{\pi_{\phi^m(K)}(\phi^{m-1}(K))}{3}.$$

Set $v(K_0) := 0$. Then $v : V \longrightarrow \mathbb{R}$ is injective, hence it induces an ordering on V . When we write down the value $v(K)$ of a ramification K in its 3-ary decomposition, we just get an algorithm, how to get from the root K_0 to K . For example 0.1121 means “go (in the driving direction) right-right-left-right”. When (ϕ, V, π) is an oriented tree with n leaves, we can extend the map ϕ to a map $\tilde{\phi} : V \setminus K_0 \cup \{1, \dots, n\} \longrightarrow V$ such that

- (i) For $1 \leq i < j \leq n$, we have $\tilde{\phi}(i) \leq \tilde{\phi}(j)$.
- (ii) For each $K \in V$, $\tilde{\phi}^{-1}(K)$ has exactly 2 elements.

The numbers $1, \dots, n$ stand for the leaves of ϕ . Furthermore, we can extend the map $v : K \longrightarrow [0, 1)$ on $\tilde{V} := V \cup \{1, \dots, n\}$ in such a way that the 3-ary decomposition of $v(i)$ describes the way from the root to the i -th leaf of ϕ , for $i = 1, \dots, n$. Then we have $v(i) < v(j)$, for $1 \leq i < j \leq n$. In consequence, we have an ordering on $\tilde{V}$.

Two trees (ϕ, V) and (ϕ', V') are called **equivalent**, if there is a bijection $f : V \longrightarrow V'$ of the ramification sets such that $f \circ \phi = \phi' \circ f$. Two oriented trees (ϕ, V, π) and (ϕ', V', π') are called **oriented equivalent**, if there is a bijection $f : V \longrightarrow V'$ of the ramification sets such that $f \circ \phi = \phi' \circ f$ and $\pi' \circ f = \pi$. When we draw oriented trees, we shall put elements K' of $\phi^{-1}(K)$ down left of K if $\pi_K(K') = 1$ and down right of K if $\pi_K(K') = 2$.

Example 2.1. The following trees with three leaves are equivalent but not oriented equivalent:



For each ramification and each leaf, we have indicated its value.


Set $\text{Ot}(n)$ to be the set of equivalence classes of oriented trees with n leaves. Observe that the cardinality $\#\text{Ot}(n)$ of $\text{Ot}(n)$ is given by the $(n-1)$ -st Catalan number $c_{n-1} = \frac{1}{n} \binom{2(n-1)}{n-1}$ which equals $\frac{(2(n-1))!}{n!(n-1)!}$. Using Stirling's

$$\sqrt{2\pi n} \cdot \left(\frac{n}{e}\right)^n \leq n! \leq 2 \cdot \sqrt{2\pi n} \cdot \left(\frac{n}{e}\right)^n,$$

we get the following:

Lemma 2.2. $\#\text{Ot}(n) < 4^{n-1}$.

The set $\text{Ot}(2)$ contains just one element. We denote it by β . The set $\text{Ot}(4)$ contains just the following elements:



For a tree (ϕ, V) and $K \in V$, there is a tree $\phi|_K$ with root K and ramifications $\{K' \in V : \phi^n(K') = K \text{ for an } n \geq 0\}$. We have to introduce several invariants: For a tree ϕ with $n > 1$ leaves and $1 \leq i \leq n$, set $w_\phi(i)$ to be the difference of the number $s_\phi(i)$ of ramifications of ϕ which are smaller than i and $i-1$. ($i-1$ is the number of leaves of ϕ , smaller than i .) For $K \in V$, set $w_\phi(K) := w_{\phi-\phi|_K}(K)$, where on the right hand-side, K is considered as leaf of $\phi - \phi|_K$.

Remark 2.3. For $K \in V$, the integer $w_\phi(K)$ is just the number of 1's arising in the 3-ary decomposition of $v(K)$.

Now, for each tree ϕ with at least 2 leaves, set $e(\phi) := (-1)^{w_\phi(1)+\dots+w_\phi(n)}$. Set $e(\tau) := 1$. For example, we have $e(\beta) = -1$. For the first tree in Example 2.1, we have $e(\phi) = -1$, and for the second tree in Example 2.1, we have $e(\phi) = +1$.

Now, let L be a graded module, ϕ an oriented tree with n leaves and $B = (b_K)_{K \in V}$ a family of bilinear maps $L \otimes L \longrightarrow L$. Recursively, we want to define a multilinear map

$$\phi(B) : L^{\otimes n} \longrightarrow L.$$

- (1) If ϕ has one leaf, i.e. B is empty, we set $\phi(B) := \text{Id}$.
- (2) If ϕ has only two leaves, i.e. $V = \{K_0\}$, for a bilinear map $b_0 : L \otimes L \longrightarrow L$, we set $\phi(b_0) := b_0$.
- (3) If $\phi^{-1}(K_0)$ contains exactly one element, say K_1 , and $\pi_{K_0}(K_1) = 1$, we set

$$\phi(B) := b_0 \circ (\phi|_{K_1}((b_K)_{K \in V \setminus K_0}) \otimes 1).$$

- (4) If $\phi^{-1}(K_0)$ contains exactly one element, say K_1 , and $\pi_{K_0}(K_1) = 2$, we set

$$\phi(B) := b_0 \circ (1 \otimes \phi|_{K_1}((b_K)_{K \in V \setminus K_0})).$$

(5) If $\phi^{-1}(K_0) = \{K_1, K_2\}$ with $\phi_{K_0}(K_1) = 1$ and $\phi_{K_0}(K_2) = 2$, we set

$$\phi(B) := b_0 \circ (\phi|_{K_1}((b_K)_{K \in V_1}) \otimes \phi|_{K_2}((b_K)_{K \in V_2})).$$

Here, V_1 denotes the ramification set of $\phi|_{K_1}$ and V_2 the ramification set of $\phi|_{K_2}$.

2.2 Operations on trees

Addition Let (ϕ, V, π) and (ϕ', V', π') be oriented trees with disjoint ramification sets. Let R be a point in neither one of them. Set $V'' := V \cup V' \cup \{R\}$. We define a map $\psi : V'' \setminus R \longrightarrow V''$ by $\psi|_{V \setminus K_0} := \phi$, $\psi|_{V' \setminus K'_0} := \phi'$ and $\psi(K_0) := \psi(K'_0) := R$.

There is a family $(\pi''_K)_{K \in W}$ of inclusions $\pi''_K : \psi^{-1}(K) \longrightarrow \{0, 1\}$ with $\pi''_K = \pi_K$, for $K \in V$, $\pi''_K = \pi'_K$ for $K \in V'$ and $\pi''_R(K_0) = 0$ and $\pi''_R(K'_0) = 1$. Now, we set

$$(\phi, V, \pi) + (\phi', V', \pi') := (\psi, V'', \pi'').$$

For example, we have $\tau + \tau = \beta$. Each tree can be reconstructed by addition out of copies of τ . It is obvious how to define the addition of non-oriented trees. The addition of oriented trees is not commutative. The addition of non-oriented trees is commutative.

Subtraction Let (ϕ, V) be a tree with n leaves and $K \in V$. Let l be the number of leaves of $\phi|_K$. The definition of a tree $\phi - \phi|_K$ with $n - l + 1$ leaves is quite obvious.

Composition Let (ϕ, V, π) be an oriented tree with n leaves and let $(\psi^{(1)}, V^{(1)}, \pi^{(1)})$, $\dots$, $(\psi^{(n)}, V^{(n)}, \pi^{(n)})$ be oriented trees. Let W be the disjoint union of V and all $V^{(i)}$. For $K \in V$ set $n(K) := 2 - |\phi^{-1}(K)|$. (This is the number of leaves belonging to K .) Let $K_1 < \dots < K_l$ all elements K of V with $n(K) > 0$. We define a map $\Phi : W \setminus K_0 \longrightarrow W$ as follows: For $K \in V \setminus K_0$, set $\Phi(K) := \phi(K)$. For $K \in V^{(i)} \setminus K_0^{(i)}$, set $\Phi(K) := \psi^{(i)}(K)$, and define the values of Φ on the $K^{(i)}$ by

$$(\Phi(K_0^{(1)}), \dots, \Phi(K_0^{(n)})) := (\underbrace{K_1, \dots, K_1}_{n_1 \text{ times}}, \dots, \underbrace{K_l, \dots, K_l}_{n_l \text{ times}}).$$

Then, (Φ, W) is a tree with a canonical orientation π' , given as follows: For each i , $K \in V^{(i)}$ and $K' \in \Phi^{-1}(K)$, we set $\pi'(K') := \pi^{(i)}(K')$. For $K \in V$ and $K' \in \Phi^{-1}(K) \cap V$, we set $\pi'(K') := \pi(K')$. It remains to define π'_{K_i} on elements of $\Phi^{-1}(K_i) \setminus V$, for $i = 1, \dots, l$. So, if $n(K_i)$ equals 2, then $\Phi^{-1}(K_i) \setminus V$ has two elements, say $K_0^{(j)}$ and $K_0^{(k)}$ with $j < k$. Set $\Phi_{K_i}(K_0^{(j)}) := 1$ and $\Phi_{K_i}(K_0^{(k)}) := 2$. If $n(K_i)$ equals 1, then $\Phi^{-1}(K_i)$ has one element in V , say K and one element which is not in V , say K' . Set $\Phi_{K_i}(K') := 1$ if $\phi_{K_i}(K) = 2$ and $\Phi_{K_i}(K') := 2$ if $\phi_{K_i}(K) = 1$.

We will denote this decomposition by $\Phi = \phi \circ (\psi^{(1)}, \dots, \psi^{(n)})$. The next lemma follows directly from the definitions:

Lemma 2.4. *In this situation, suppose that there is a family $B = (b_K)_{K \in W}$ of bilinear maps $L \otimes L \longrightarrow L$. Set $B^{(0)} := (b_K)_{K \in V}$ and $B^{(i)} := (b_K)_{K \in V^{(i)}}$, for $i = 1, \dots, n$. We have*

$$\phi \circ (\psi^{(1)}, \dots, \psi^{(n)})(B) = (-1)^{\text{exponent}} \phi(B^{(0)}) \circ (\psi^{(1)}(B^{(1)}) \otimes \dots \otimes \psi^{(n)}(B^{(n)})),$$

where the exponent is the sum

$$\left(\sum_{K \in V^{(1)}} b_K \right) \left(\sum_{K \in V}^{V > 1} b_K \right) + \dots + \left(\sum_{K \in V^{(n-1)}} b_K \right) \left(\sum_{K \in V}^{V > n-1} b_K \right).$$

We remind that $V > i$ means that the value $v(V)$ is greater than the value $v(i)$ of the i -th leaf of ϕ .

3 Decomposition of differential graded Lie algebras

Let $L = (L, d, [\cdot, \cdot])$ be a differential graded Lie algebra, where the differential d is of degree $+1$. Suppose that there is a splitting η , i.e. a map of degree -1 such that $d\eta d = d$. Furthermore, suppose that $\eta^2 = 0$ and $\eta d\eta = \eta$. When we use a Lie bracket on $\text{Hom}(L, L)$, we mean the graded commutator. Observe that $[d, \eta]$ is a projector, and we can identify the cohomology $H(L)$ with the graded submodule $H := \text{Fix}(1 - [d, \eta])$ of L . Set F to be the d -invariant submodule $\text{Fix}([d, \eta])$ of L . We can consider the pair (F, d) as an L_∞ -algebra. As DG module, we have a decomposition $(L, d) \cong (H, 0) \oplus (F, d)$. The aim of this section is to construct a minimal L_∞ -structure on H such that the decomposition $L \cong H \oplus F$ still holds in the category of L_∞ -algebras. We begin with the construction of an L_∞ -algebra map $g : F \rightarrow L$.

In the next theorem, for a tree $\phi \in \text{Ot}(n)$, we write Φ for the multilinear map $\phi([\cdot, \cdot], \dots, [\cdot, \cdot]) : L^{\otimes n} \longrightarrow L$.

Theorem 3.1. *The following graded anti-symmetric maps $g_n : F^{\otimes n} \longrightarrow L$ of degree $1 - n$ define a morphism $g : F \longrightarrow L$ of L_∞ -algebras:*

$$\begin{aligned} g_1 &:= \text{inclusion} \\ g_2 &:= \frac{-1}{2} \cdot [\eta \cdot, \cdot] \circ \alpha_2 \\ &\vdots \\ g_n &:= \left(\frac{-1}{2}\right)^{n-1} \sum_{\phi \in \text{Ot}(n)} \Phi \circ (\eta^{n-1} \otimes 1) \circ \alpha_n \end{aligned}$$

Proof.

$$\begin{aligned}
& (-1)^{n-1} g_n \circ (d \otimes 1^{\otimes n-1}) \circ \alpha_{1,n} = \\
& \left(\frac{1}{2}\right)^{n-1} \sum_{\phi \in \text{Ot}(n)} \sum_{i+j=n-1} \Phi \circ (\eta^{\otimes n-1} \otimes 1) \circ (1^{\otimes i} \otimes d \otimes 1^{\otimes j}) \circ \alpha_n = \\
& \left(\frac{-1}{2}\right)^{n-1} \sum_{\phi \in \text{Ot}(n)} \sum_{i+j=n-2} (-1)^{i-1} \cdot \Phi \circ (\eta^{\otimes i} \otimes 1 \otimes \eta^{\otimes j} \otimes 1) \circ \alpha_n + \\
& \left(\frac{-1}{2}\right)^{n-1} \sum_{\phi \in \text{Ot}(n)} \sum_{i+j=n-1} \Phi \circ (1^{\otimes i} \otimes d \otimes 1^{\otimes j}) \circ (\eta^{\otimes n-1} \otimes 1) \circ \alpha_n = \\
& \left(\frac{-1}{2}\right)^{n-1} \sum_{i+j=n} (-1)^i \sum_{\phi_1 \in \text{Ot}(i)} \sum_{\phi_2 \in \text{Ot}(j)} [\Phi_1 \circ (\eta^{i-1} \otimes 1) \circ \alpha_i, \Phi_2 \circ (\eta^{j-1} \otimes 1) \circ \alpha_j] \circ \alpha_{i,n} + \\
& \left(\frac{-1}{2}\right)^{n-1} \sum_{\phi \in \text{Ot}(n)} d \circ \Phi \circ (\eta^{\otimes n-1} \otimes 1) \circ \alpha_n = \\
& \frac{-1}{2} \sum_{i+j=n} (-1)^i [g_i, g_j] \circ \alpha_{i,n} + d g_n.
\end{aligned}$$

Thus, condition (1.7) is verified. $\square$


Remark that in the proof of Theorem 3.1, we didn't make use of the Jacobi identity. However, the Jacobi identity effects that many of the terms in the decomposition of g_n annihilate each other. Before using the formulas in a computer algorithm, one has to study this more closely.

Now we come to the slightly more difficult construction of an L_∞ -algebra structure μ_* on $H := H(L, d)$ with $\mu_1 = 0$ and of an L_∞ -quasi-isomorphism $f : H \rightarrow L$. In the A_∞ -context, the existence of an A_∞ -structure on the cohomology of a DG algebra A had already be shown (for A connected) by Kadeishvili [6], Gugenheim/ Stasheff [3] and (in the general case) by Merkulov [16]. Merkulov gives a recursion formula for construction of the higher products. A similar recursion formula for the L_∞ -case can be found in the article [4] of Huebschmann and Stasheff. Kontsevich and Soibelman [9] rewrote the higher terms obtained by Merkulov's construction (A_∞ -case) in terms of decorated trees. Their formulas are still recursion formulas.

We have to make some preparations. First of all, there is the following simple but important lemma:

Lemma 3.2. *Let $n \geq 3$ be a natural number. There is a 1:1-correspondence between triples (Φ, K, σ) , where $\Phi = (\Phi, V, \pi)$ is an oriented tree with n leaves, K is a ramification in V , σ a permutation in Σ_n and 6-tuples $(k, \phi, \psi, \rho, \gamma, \delta)$, where k is a natural number with $2 \leq k \leq n-1$, ϕ is a tree in $\text{Ot}(k)$, ψ is a tree in $\text{Ot}(l)$ where*

$l := n + 1 - k$, ρ is a shuffle in $\text{Sh}(k, n)$ and $\gamma \in \Sigma_l$, $\delta \in \Sigma_k$ are permutations.



Example:

The fine lines represent the tree ψ and the fat lines the tree ϕ .

In the sequel, the first r leaves of Φ will be associated to the indexes $\sigma(1) = \rho(\gamma(1) + k - 1), \dots, \sigma(r) = \rho(\gamma(r) + k - 1)$, the following k leaves to the indexes $\sigma(r + 1) = \rho(\delta(1)), \dots, \sigma(r + k) = \rho(\delta(k))$ and the remaining leaves to the indexes $\sigma(r + k + 1) = \rho(\gamma(r + 2) + k - 1), \dots, \sigma(n) = \rho(\gamma(l) + k - 1)$.

To the triple (Φ, K, σ) , we associate the following data: Set k to be the number of leaves of $\Phi|_K$, $\phi := \Phi|_K$, $\psi := \Phi - \phi$. Let r be the number of leaves F of Φ with $F < K$. The shuffle ρ is chosen in such a way that $\{\rho(1), \dots, \rho(k)\} = \{\sigma(r + 1), \dots, \sigma(r + k)\}$. The permutation δ is defined by $\delta(i) := \rho^{-1}(\sigma(r + i))$, for $i = 1, \dots, k$ and γ is defined in the following way:

$$\gamma(i) := \begin{cases} \rho^{-1}(\sigma(i)) - k + 1 & \text{for } i = 1, \dots, r \\ 1 & \text{for } i = r + 1 \\ \rho^{-1}(\sigma(i + k - 1)) - k + 1 & \text{for } i = r + 2, \dots, l \end{cases}$$

In the other direction, to the 6-tuple $(k, \phi, \psi, \rho, \gamma, \delta)$, we associate the following data: Set $r := \gamma^{-1}(1) - 1$. Then Φ is the composition

$$\Phi = \psi \circ (\underbrace{\tau, \dots, \tau}_{r \text{ times}}, \phi, \tau, \dots, \tau),$$

where τ again stands for the tree with one leaf. The ramification K is the root of ϕ , considered as ramification of Φ and σ is given by

$$\sigma(i) := \begin{cases} \rho(\gamma(i) + k - 1) & \text{for } i = 1, \dots, r \\ \rho(\delta(i - r)) & \text{for } i = r + 1, \dots, r + k \\ \rho(\gamma(i - (k - 1)) + k - 1) & \text{for } i = r + k + 1, \dots, n. \end{cases}$$

Now suppose that such corresponding tuples $(\Phi, \hat{K}, \sigma)$ and $(k, \phi, \psi, \rho, \gamma, \delta)$ are given. Let V' be the ramification set of ψ and V'' the ramification set of ϕ . Then $V := V' \cup V''$ is the ramification set of Φ . Again, set $r := \gamma^{-1}(1) - 1$. Remark that the ordering on V depends on γ . We define a permutation $\tilde{\gamma} \in \Sigma_{l-1}$ by

$$\tilde{\gamma}(i) := \begin{cases} \gamma(i) - 1 & \text{for } i = 1, \dots, r \\ \gamma(i + 1) - 1 & \text{for } i = r + 1, \dots, l - 1. \end{cases}$$

Lemma 3.3. We keep all notation from above. Let $B = (b_K)_{K \in V}$ be a family of homogeneous bilinear forms $L \otimes L \rightarrow L$. Denote the subfamilies $(b_K)_{K \in V'}$ and

$(b_K)_{K \in V''}$ by B' and B'' . Set W to be the set of all ramifications $K \in V$ such that $K > \hat{K}$. Then we have

$$\begin{aligned} & \psi(B') \circ \gamma \circ (\phi(B'') \circ \delta \otimes 1 \otimes \dots \otimes 1) \circ \rho \\ &= (-1)^{r+rk} \psi(B') \circ \underbrace{(1 \otimes \dots \otimes 1)}_{r \text{ times}} \otimes \phi(B'') \otimes 1 \otimes \dots \otimes 1 \circ \sigma \\ &= (-1)^{r+rk+\sum_{K \in W} b_K \cdot B''} \Phi(B) \circ \sigma. \end{aligned}$$

Proof. Let $a_1, \dots, a_n$ be homogeneous elements of L . We get

$$\begin{aligned} & (\psi(B') \circ \gamma \circ (\phi(B'') \circ \delta \otimes 1 \otimes \dots \otimes 1) \circ \rho)(a_1 \otimes \dots \otimes a_n) = \\ & \chi(\rho, a_1, \dots, a_n) \chi(\delta, a_{\rho(1)}, \dots, a_{\rho(k)}). \\ & (\psi(B') \circ \gamma) \underbrace{(\phi(B'')(a_{\rho(\delta(1))}, \dots, a_{\rho(\delta(k))}))}_{=: u_1} \otimes \underbrace{a_{\rho(k+1)}}_{u_2} \otimes \dots \otimes \underbrace{a_{\rho(n)}}_{u_l} = \\ & \chi(\rho, a_1, \dots, a_n) \chi(\delta, a_{\rho(1)}, \dots, a_{\rho(k)}) \chi(\gamma, u_1, \dots, u_l) \psi(B')(u_{\gamma(1)}, \dots, u_{\gamma(l)}). \end{aligned}$$

Using the following three formulas

$$\begin{aligned} & \chi(\sigma, a_1, \dots, a_n) = (-1)^{kr+(a_{\sigma(1)}+\dots+a_{\sigma(r)})(a_{\rho(1)}+\dots+a_{\rho(k)})} \chi(\rho, a_1, \dots, a_n). \\ & \chi(\delta, a_{\rho(1)}, \dots, a_{\rho(k)}) \chi(\tilde{\gamma}, a_{\rho(k+1)}, \dots, a_{\rho(n)}), \\ & u_{\gamma(1)} \otimes \dots \otimes u_{\gamma(l)} = (-1)^{B''(a_{\sigma(1)}+\dots+a_{\sigma(r)})}. \\ & (1 \otimes \dots \otimes 1 \otimes \phi(B'') \otimes 1 \otimes \dots \otimes 1)(a_{\sigma(1)} \otimes \dots \otimes a_{\sigma(n)}), \\ & \chi(\gamma, u_1, \dots, u_l) = (-1)^{r+u_1(u_{\gamma(1)}+\dots+u_{\gamma(r)})} \chi(\tilde{\gamma}, u_2, \dots, u_l), \end{aligned}$$

this expression is just

$$(-1)^{kr+r} \chi(\sigma, a_1, \dots, a_n) \psi(B')((1 \otimes \dots \otimes 1 \otimes \phi(B'') \otimes 1 \otimes \dots \otimes 1)(a_{\sigma(1)} \otimes \dots \otimes a_{\sigma(n)})).$$

The second equality of this Lemma is just a special case of Lemma 2.4. $\square$

We turn to the construction of an L_∞ -structure on $H(L)$.

Proposition 3.4. *The map $[d, \eta] = d\eta + \eta d$ is a projection, i.e. $[d, \eta]^2 = [d, \eta]$. And $H := \text{Kern}[d, \eta]$ is as module isomorphic to $H(L)$. Remark that under the assumption of the beginning of this section, we have*

$$H = \text{Kern } d \cap \text{Kern } \eta.$$

The bracket on L induces a Lie-bracket on $H(L)$ and the induced bracket on H (via the isomorphism $H \longrightarrow H(L)$) is just given by $(1 - d\eta)[\cdot, \cdot] = (1 - [d, \eta])[\cdot, \cdot]$.

For simplicity, we set $g := \eta[\cdot, \cdot]$.

Theorem 3.5. *The following graded anti-symmetric maps $\mu_n : H^{\otimes n} \longrightarrow H$ of degree $2 - n$ define the structure of an L_∞ -algebra on H :*

$$\begin{aligned}\mu_1 &:= 0 \\ \mu_2 &:= (1 - d\eta)[\cdot, \cdot] \\ &\vdots \\ \mu_n &:= \left(\frac{-1}{2}\right)^{n-1} \sum_{\phi \in \text{Ot}_n} e(\phi) \phi((1 - [d, \eta])[\cdot, \cdot], g, \dots, g) \circ \alpha_n\end{aligned}$$

Here, the sum is taken over all trees ϕ with n leaves and $\phi((1 - [d, \eta])[\cdot, \cdot], g, \dots, g)$ is the n -linear form obtained by assigning the bilinear form $(1 - [d, \eta])[\cdot, \cdot]$ to the root of the tree ϕ and the bilinear form g to each other ramification. The sign $e(\phi)$ is defined in Section 2. and α_n is the anti-symmetrization map.

Proof. We must show that

$$\sum_{k+l=n+1} (-1)^{k(l-1)} \mu_l \circ (\mu_k \otimes 1 \otimes \dots \otimes 1) \circ \alpha_{k,n} = 0. \quad (3.8)$$

Up to the factor $(-1)^{n-1}$ this sum has the form

$$\begin{aligned}\sum_k \sum_{\phi, \psi} \sum_{\rho, \gamma, \delta} (-1)^{k(l-1)} e(\phi) e(\psi) \psi((1 - [d, \eta])[\cdot, \cdot], g, \dots, g) \circ \gamma \circ \\ \circ (\phi((1 - [d, \eta])[\cdot, \cdot], g, \dots, g) \circ \delta \otimes 1 \otimes \dots \otimes 1) \circ \rho, \quad (3.9)\end{aligned}$$

where k ranges from 2 to $n-1$, $l = n+1-k$, ϕ and ψ vary in $\text{Ot}(k)$ and $\text{Ot}(l)$, ρ in $\text{Sh}(k, n)$, γ and δ in Σ_l and Σ_k . For corresponding tuples $(k, \phi, \psi, \rho, \gamma, \delta)$ and $(\Phi, \hat{K}, \sigma)$ as in Lemma 3.2, we denote as usual $r := \gamma^{-1}(1) - 1$ and by t the number of ramifications of ψ , greater than $r+1$. Using

$$\begin{aligned}e(\Phi) &= (-1)^{w_\Phi(\hat{K})(k-1)} e(\phi) e(\psi) \\ w_\Phi(\hat{K}) &= l - 1 - r - t\end{aligned}$$

and Lemma 3.3, the expression (3.9) can be expressed as

$$\sum_{\Phi \in \text{Ot}(n)} \sum_{\hat{K} \in V \setminus K_0} e(\Phi) (-1)^{r+w_\Phi(\hat{K})} \Phi(B) \circ \alpha_n,$$

where $B = (B_K)_{K \in V}$ is the family with $b_{K_0} = b_{\hat{K}} = (1 - [d, \eta])[\cdot, \cdot]$ and $b_K = \eta[\cdot, \cdot]$ for $K \neq K_0, \hat{K}$. To show that the last term is zero, it is enough to show the following two conditions:

$$\sum_{\Phi \in \text{Ot}(n)} \sum_{K \in V \setminus K_0} \sum_{\sigma \in \Sigma_n} (-1)^{r+w_\Phi(K)} e(\Phi) \Phi((1 - [d, \eta])[\cdot, \cdot], g, \dots, g, \underbrace{[\cdot, \cdot]}_{\text{pos. } K}, g, \dots, g) \circ \sigma = 0. \quad (3.10)$$

For each tree Φ , we have

$$\sum_{K \in V \setminus K_0} (-1)^{r+w_\Phi(K)} e(\Phi) \Phi((1 - [d, \eta])([\cdot, \cdot], g, \dots, g, \underbrace{[d, \eta](\cdot, \cdot)}_{\text{position } K}, g, \dots, g) \circ \sigma = 0. \quad (3.11)$$

The first condition follows by the Jacobi-identity and an easy combinatorial argument. In equation (3.11) the term annihilate each other since the differential d trickles down the branches of Φ :

Initiation of the trickling: Suppose that $\Phi^{-1}(K_0)$ contains an element K' with $\pi_{K_0}(K') = 1$. We have the following picture:

$$0 = \begin{array}{c} (1 - [d, \eta])d[\cdot, \cdot] \\ \swarrow \quad \searrow \\ \eta[\cdot, \cdot] \quad \eta[\cdot, \cdot] \end{array} = \begin{array}{c} (1 - [d, \eta])([\cdot, \cdot] \\ \swarrow \quad \searrow \\ d\eta[\cdot, \cdot] \quad \eta[\cdot, \cdot] \end{array} + (-1)^{\# \text{ramific. of } \Phi|_{K'}} \begin{array}{c} (1 - [d, \eta])([\cdot, \cdot] \\ \swarrow \quad \searrow \\ \eta[\cdot, \cdot] \quad d\eta[\cdot, \cdot] \end{array}$$

Here, we only have drawn the top of the tree Φ for the case where $\Phi^{-1}(K_0)$ consists of two elements K', K'' and the corresponding bilinear forms. It is quite obvious how this goes when $\Phi^{-1}(K_0)$ has only one element, since $d|_H = 0$.

Going-on of the trickling at a ramification $K \in V$: We illustrate the case, where $\Phi^{-1}(K)$ has two elements K', K'' with $\pi_K(K') = 1$.

$$\begin{array}{c} \eta d[\cdot, \cdot] \\ \swarrow \quad \searrow \\ \eta[\cdot, \cdot] \quad \eta[\cdot, \cdot] \end{array} - \begin{array}{c} \eta[\cdot, \cdot] \\ \swarrow \quad \searrow \\ d\eta[\cdot, \cdot] \quad \eta[\cdot, \cdot] \end{array} - (-1)^{\# \text{ramific. of } \Phi|_{K'}} \begin{array}{c} \eta[\cdot, \cdot] \\ \swarrow \quad \searrow \\ \eta[\cdot, \cdot] \quad d\eta[\cdot, \cdot] \end{array} = 0$$

Iterating the trickling down to the leaves and using $d|_H = 0$, we see that all terms in the sum are annihilated. $\square$

Remark 3.6. The restriction of μ defines a formal map $H^1 \longrightarrow H^2$. One can show that this is just the Kuranishi map as defined in [10].

Theorem 3.7. *The following anti-symmetric maps $f_n : H^{\otimes n} \longrightarrow L$ of degree $1 - n$ define an L_∞ -equivalence $H \longrightarrow L$ (i.e. an L_∞ -quasi-isomorphism).*

$$\begin{aligned} f_1 &:= \text{inclusion} \\ f_2 &:= -g \\ &\vdots \\ f_n &:= -\left(\frac{-1}{2}\right)^{n-1} \sum_{\phi \in \text{Ot}(n)} e(\phi) \phi(g, \dots, g) \circ \alpha_n. \end{aligned}$$

Proof. For $n \geq 0$, we have to prove the equation

$$df_n - \sum_{i+j=n} \frac{(-1)^i}{2} [f_i, f_j] \alpha_{i,n} = \sum_{k+l=n+1} (-1)^{k(l-1)} f_l \circ (\mu_k \otimes 1 \otimes \dots \otimes 1) \circ \alpha_{k,n}$$

For $l = 1$, the right hand-side is just μ_n . Since

$$df_n = \left(\frac{-1}{2}\right)^{n-1} \sum_{\phi \in \text{Ot}(n)} e(\phi) \phi(-d\eta[\cdot, \cdot], g, \dots, g) \circ \alpha_n,$$

it is sufficient to show the following three identities:

$$- \sum_{i+j=n} \frac{(-1)^i}{2} [f_i, f_j] \alpha_{i,n} = \left(\frac{-1}{2}\right)^{n-1} \sum_{\phi \in \text{Ot}(n)} e(\phi) \phi([\cdot, \cdot], g, \dots, g) \circ \alpha_n \quad (3.12)$$

$$f_l \circ (\phi([\cdot, \cdot], g, \dots, g) \circ \alpha_k \otimes 1 \otimes \dots \otimes 1) \circ \alpha_{k,n} = 0 \text{ for } l > 1, k + l = n + 1. \quad (3.13)$$

$$\begin{aligned} & \left(\frac{-1}{2}\right)^{n-1} \sum_{\phi \in \text{Ot}(n)} e(\phi) \phi(\eta d[\cdot, \cdot], g, \dots, g) \circ \alpha_n = \\ & - \sum_{\substack{k+l=n+1 \\ l \geq 2}} (-1)^{k(l-1)} \sum_{\phi \in \text{Ot}(k)} \left(\frac{-1}{2}\right)^{k-1} e(\phi) \cdot \\ & f_l \circ (\phi(d\eta + \eta d)[\cdot, \cdot], g, \dots, g) \circ \alpha_k \otimes 1 \otimes \dots \otimes 1) \circ \alpha_{k,n}. \end{aligned} \quad (3.14)$$

Proof of equation (3.14): The right hand-side of equation (3.14) is

$$\begin{aligned} & \left(\frac{-1}{2}\right)^{n-1} \sum_{k+l=n+1} \sum_{\phi, \psi} \sum_{\gamma, \delta, \rho}^{k, l \geq 2} (-1)^{k(l-1)} e(\phi) e(\psi) \psi(g, \dots, g) \circ \gamma \circ \\ & (\phi([d, \eta][\cdot, \cdot], g, \dots, g) \circ \delta \otimes 1 \otimes \dots \otimes 1) \circ \rho. \end{aligned}$$

As in the proof of Theorem 3.5, this expression takes the form

$$\left(\frac{-1}{2}\right)^{n-1} \sum_{\Phi \in \text{Ot}(n)} \sum_{\hat{K} \in V} \sum_{\sigma \in \Sigma_n} (-1)^{r+w_{\Phi}(\hat{K})} e(\Phi) \Phi(B) \sigma,$$

where $B = (b_K)_{K \in V}$ is the family with $b_{\hat{K}} = [d, \eta][\cdot, \cdot]$ and $b_K = \eta[\cdot, \cdot]$ for $K \neq \hat{K}$.

Hence to show equation (3.14), it is enough to show that for each tree ϕ , we have

$$\Phi(\eta d[\cdot, \cdot], g, \dots, g) = \sum_{K \in V \setminus K_0} (-1)^{r+w_{\Phi}(K)} \Phi(B).$$

This is true by the same trickling argument as in Theorem 3.5.

Proof of equation (3.13): This is again the Jacobi-identity and some combinatorics.

Proof of equation (3.12):

$$\begin{aligned}
\sum_{i+j=n} \frac{(-1)^i}{2} [\cdot, \cdot] \circ (f_i \otimes f_j) \circ \alpha_{i,n} &= \\
= \sum_{i+j=n} \left(-\frac{1}{2}\right)^{i-1+j-1} \sum_{\phi \in \text{Ot}(i), \psi \in \text{Ot}(j)} \frac{(-1)^i}{2} e(\phi)e(\psi)(\phi + \psi)([\cdot, \cdot], g, \dots, g) \circ \alpha_n \\
&= -\left(-\frac{1}{2}\right)^{n-1} \sum_{\Phi \in \text{Ot}(n)} e(\Phi)\Phi([\cdot, \cdot], g, \dots, g) \circ \alpha_n.
\end{aligned}$$

□

Almost in the same manner, one can realize the following: If $L = (L, d, [\cdot, \cdot])$ is a DG Lie algebra and $f_1 : (M, d^M) \longrightarrow (L, d)$ a homotopy equivalence between DG modules, then there is an L_∞ -algebra structure μ_* on M with $\mu_1 = d^M$ and an L_∞ -quasi-isomorphism $f : (M, \mu_*) \longrightarrow (L, d, [\cdot, \cdot])$, extending f_1 . In other words: An up to homotopy differential graded Lie algebra is an L_∞ -algebra. In the A_∞ -context, this was already shown by Markl [14]. Applying Proposition 1.2 to the direct sum $f \oplus g$, we get the decomposition theorem for differential graded Lie algebras:

Theorem 3.8. *The direct sum*

$$f \oplus g : (H, \mu) \oplus (F, d) \longrightarrow (L, d, [\cdot, \cdot])$$

is an isomorphism of L_∞ -algebras.

One can show less constructively but more generally (see [7]) that each L_∞ -algebra with splitting is isomorphic to the direct sum of a minimal and a linear contractible one.

Corollary 3.9. *If $(L, d, [\cdot, \cdot])$ and $(L', d', [\cdot, \cdot])$ are DG Lie algebras such that (L, d) and (L', d') are split, then, for each L_∞ -quasi-isomorphism $f : L \longrightarrow L'$, there exists an L_∞ -morphism $g : L' \longrightarrow L$ such that f_1 and g_1 are inverse maps on the homology.*

4 Palamodov DG manifolds

4.1 Palamodov spaces

We want to study a class of DG manifolds (M, Q^M) , for which the vectorfield Q^M is not only formal but satisfies a convergence condition. Thus, the underlying graded vector space M should be a normed or at least a metric vector space. It is natural, for instance, to define “Banach DG manifolds”, where M is a graded Banach space and the formal sum $\sum_n \frac{1}{n!} \tilde{Q}^M$ converges in a neighbourhood of $0 \in M$. But in view

of applications in analytic deformation theory, this is not general enough since, for example the tangent complex of a complex space is not Banach. We have to work more generally with Palamodov's *RO*- (or in Cyrillic *PO*-) *spaces*, whose topology is defined by a countable family of semi-norms. We shall call these spaces *Palamodov spaces* instead of *RO-spaces*. We will define *Palamodov DG manifolds* and correspondingly *Palamodov L_∞ -algebras*. This generalises Bingener and Kosarew's notion [2] of *graded PO Lie algebras*.

Definition 4.1. A **Palamodov space** is a vectorspace E together with a family $(\|\cdot\|_\lambda)_{0 < \lambda < 1}$ of semi-norms such that

$$\|\cdot\|_\lambda \leq \|\cdot\|_{\lambda'} \quad \text{for} \quad \lambda < \lambda'.$$

A morphism $f : E \longrightarrow F$ of Palamodov spaces is a morphism of vectorspaces together with global constants $\epsilon \in (0, 1)$ and $C > 0$ such that, for each element $e \in E$ and each $\lambda \in (0, 1)$, we have

$$\|f(e)\|_\lambda \leq C \cdot \|e\|_\lambda.$$

Example 4.2. If E is the space of holomorphic functions on an open polydisk P of poly-radius $r \in \mathbb{R}_{>0}^N$ we can define seminorms

$$|f|_\lambda := \sup\{|f(z)| : z \in \lambda \cdot P\},$$

for $0 < \lambda < 1$. By Cauchy's integral formula, we have the estimation

$$(\lambda' - \lambda)^2 \cdot |\partial f / \partial x_i|_\lambda \leq |f|_{\lambda'},$$

for $\lambda < \lambda' < 1$. If we set

$$\|\sum_\nu a_\nu x^\nu\|_\lambda := \sum_\nu |a_\nu| \cdot |x^\nu|_\lambda = \sum_\nu |a_\nu| \cdot \lambda^{|\nu|} \cdot r^\nu,$$

and $C := \frac{1}{\epsilon \cdot r_i}$, we get the better estimation

$$(\lambda' - \lambda) \cdot \|\partial f / \partial x_i\|_\lambda \leq \frac{C}{\epsilon} \cdot \|f\|_{\lambda'},$$

for $1 - \epsilon < \lambda < \lambda' < 1$. For the following reason, we usually use the Palamodov structure given by $\|\cdot\|_\lambda$ on E : Let $R^0 = \Gamma(P, \mathcal{O}_P)$ be the $\mathbb{C}$ -algebra of holomorphic functions on the open polydisk P and $R = R^0[x_i]_{i \in I}$ a finitely generated free DG algebra over R^0 with $g(x_i) < 0$, for all $i \in I$. The graded vector space $L = \text{Der}(R, R)$ is then a sum of copies of E and carries the structure of a Palamodov DG Lie algebra (see the definition below).

Multilinear maps and power series In this section, E and F are graded vector spaces. By definition, a **polynomial $E \rightarrow F$ of (polynomial) degree p and linear degree i** is a map of the form $x \mapsto \tilde{\phi}(x)$, where $\phi : E^{\odot p} \rightarrow F$ is a *graded symmetric* p -linear map (of linear degree i) and $\tilde{\phi}(x) = \phi(x, \dots, x)$. Following Bingener and Kosarew [2], for Palamodov spaces E and F and a given $\epsilon \in (0, 1)$, we define the following pseudo-norms on the graded vectorspace of polynomial maps $u : E \rightarrow F$ of polynomial degree p :

$$\|u\|^{0,\epsilon} := \sup\{\|u(x)\|_{\lambda} : \|x\|_{\lambda} \leq 1 \text{ and } 1 - \epsilon \leq \lambda < 1\}$$

$$\|u\|^{1,\epsilon} := \sup\{(\lambda' - \lambda)^{p-1} \|u(x)\|_{\lambda} : \|x\|_{\lambda'} \leq 1 \text{ and } 1 - \epsilon \leq \lambda < \lambda' < 1\}$$

Furthermore, we define the following pseudo-norms on the graded vectorspace of p -linear maps $\phi : E^{\otimes p} \rightarrow F$.

$$|\phi|^{0,\epsilon} := \sup\{\|\phi(x_1, \dots, x_p)\|_{\lambda} : \|x_i\|_{\lambda} \leq 1 \quad \forall i \text{ and } 1 - \epsilon \leq \lambda < 1\}$$

$$|\phi|^{1,\epsilon} := \sup\{(\lambda' - \lambda)^{p-1} \|\phi(x_1, \dots, x_p)\|_{\lambda} : \|x_i\|_{\lambda'} \leq 1 \quad \forall i \text{ and } 1 - \epsilon \leq \lambda < \lambda' < 1\}$$

Remark 4.3. Suppose that E and F are graded Palamodov spaces. For each graded symmetric p -linear map $\phi : E^{\odot p} \rightarrow F$, we have

$$\|\phi\|^{1,\epsilon} \leq |\phi|^{1,\epsilon} \leq \frac{p^p}{p!} \|\tilde{\phi}\|^{1,\epsilon}.$$

For graded vector spaces E and F , set $F[[E]]^i$ to be the set of formal sums $\sum_{p=0}^{\infty} u_p$, where $u_p : E \rightarrow F$ is a polynomial of degree p and linear degree i . By omission, we mean $i = 0$ or another fixed $i \in \mathbb{Z}$. A formal map $u = \sum_{p=0}^{\infty} u_p$ in $F[[E]]$ is called **strictly convergent** (with respect to ϵ), if there exists a $t > 0$ such that

$$\sum_{p=0}^{\infty} \|u_p\|^{0,\epsilon} \cdot t^p < \infty.$$

A power series $u = \sum_{p=0}^{\infty} u_p$ in $F[[E]]$ is called **1-convergent** (with respect to ϵ), if there exists a $t > 0$ such that

$$\sum_{p=0}^{\infty} \|u_p\|^{1,\epsilon} \cdot t^p < \infty.$$

Each strictly convergent formal map is 1-convergent.

Palamodov DG manifolds

Definition 4.4. For $i \in \{-max, \dots, +max\}$, let M^i be Palamodov spaces and $M := \oplus M^i$. A coderivation $Q \in \text{Coder}^{\geq 0}(SM)$ on SM is called **1-convergent** (resp. **strictly convergent**) with respect to ϵ , if the formal sum $\tilde{Q} := \sum_n \frac{1}{n!} \tilde{Q}_n$ in $M[[M]]^1$ is 1-convergent (resp. strictly convergent) with respect to ϵ . A coderivation Q is

called **convergent** with respect to ϵ , if for each component $Q_n \in \text{Mult}_n(M, M)$, we have $|Q_n|^{1,\epsilon} < \infty$ and if the formal sum $\tilde{Q} := \sum_n \frac{1}{n!} \tilde{Q}_n$, considered as formal map $M^{\geq 0} \longrightarrow M^{\geq 0}$ is strictly convergent with respect to ϵ . Denote the set of all convergent coderivations of SM of degree i by $\text{Coder}^i(SM)$ and set $\text{Coder}(SM) := \oplus_{i \geq 0} \text{Coder}^i(SM)$.

We want to emphasise that by graded symmetry, the odd components of m only give a linear contribution to the term $\tilde{Q}(m)$.

Proposition 4.5. *The sets of 1-convergent, strictly convergent and convergent coderivations, respectively of SM are graded Lie subalgebras of $\text{Coder}^{\geq 0}(SM)$.*

Proof. We only prove the statement for 1-convergent coderivations. We must show that, if Q and q are 1-convergent coderivations, then $[Q, q]$ is 1-convergent. Fix λ_0, λ_1 such that $1 - \epsilon \leq \lambda_0 < \lambda_1 < 1$. Set $\lambda := \frac{1}{2}(\lambda_1 - \lambda_0)$.

$$\begin{aligned} & (\lambda_1 - \lambda_0)^{n-1} ||[Q, q]_n^\sim||_{\lambda_0} \leq \\ & 2^{n-1} (\lambda - \lambda_0)^{n-1} \sum_{k+l=n+1} (||Q_l(\tilde{q}_k(x), x, \dots, x)||_{\lambda_0} + ||q_l(\tilde{Q}_k(x), x, \dots, x)||_{\lambda_0}) \leq \\ & 2^{n-1} (\lambda_1 - \lambda)^k \sum_{k+l=n+1} (||Q_l|^{1,\epsilon}||\tilde{q}_k(x)||_\lambda ||x||_\lambda^{l-1} + ||q_l|^{1,\epsilon}||\tilde{Q}_k(x)||_\lambda ||x||_\lambda^{l-1}) \leq \\ & 2^{n-1} \sum_{k+l=n+1} (||Q_l|^{1,\epsilon}||q_k||^{1,\epsilon} + ||q_l|^{1,\epsilon}||Q_k||^{1,\epsilon}) ||x||_{\lambda_1}^n \leq \\ & \frac{2^{n-1} (n-1)^{n-1}}{(n-1)!} \sum (||Q_l|^{1,\epsilon}||q_k||^{1,\epsilon} + ||q_l|^{1,\epsilon}||Q_k||^{1,\epsilon}) ||x||_{\lambda_1}^n \end{aligned}$$

Thus,

$$||[Q, q]_n^\sim||^{1,\epsilon} \leq \frac{2^{n-1} (n-1)^{n-1}}{(n-1)!} \sum (||Q_l|^{1,\epsilon}||q_k||^{1,\epsilon} + ||q_l|^{1,\epsilon}||Q_k||^{1,\epsilon}).$$

By the Stirling formula, we have $\frac{2^{n-1} (n-1)^{n-1}}{(n-1)!} \leq (2e)^{n-1}$, hence

$$\sum_n \frac{1}{n!} ||[Q, q]_n^\sim||^{1,\epsilon} t^n \leq 2 \left(\sum_k \frac{(2e)^k}{k!} ||\tilde{Q}_k||^{1,\epsilon} t^k \right) \left(\sum_l \frac{(2e)^l}{l!} ||\tilde{q}_l||^{1,\epsilon} t^l \right),$$

which is finite, for small t . □

Definition 4.6. A formal DG manifold (M, Q^M) , where $M = \oplus_{i=-d}^d M^i$ is a graded Palamodov space and $Q^M \in \text{Coder}^1(SM)$ will be called **Palamodov DG manifold**. If M is Banach and the Palamodov structure is given by $|| \cdot ||_\lambda := || \cdot ||$, for each $\lambda \in (0, 1)$, then (M, Q^M) will be called **Banach DG manifold**. A **morphism** $f : M \longrightarrow N$ of Palamodov DG manifolds is a morphism f of formal DG manifolds such that $||f_n||^{1,\epsilon}$ is finite, for each n , and such that the formal map $\tilde{f} = \sum_n \frac{1}{n!} \tilde{f}_n$ is strictly convergent as a map $M^{\geq 0} \longrightarrow M^{\geq 0}$.

The following inverse function theorem shows that Palamodov DG manifolds are well behaved:

Proposition 4.7. *If $f : M \rightarrow N$ is a morphism of Palamodov super manifolds (resp. Palamodov DG manifolds) such that f_1 is an isomorphism of Palamodov spaces, then f is an isomorphism.*

Proof. The proof of Proposition II.8.4 of [2] applies directly to this situation. For convenience, we indicate it, however. It uses the following statement: Let $\sum_p \gamma_p t^p$ the Taylor series at zero of the function $\frac{1}{2} - \frac{1}{4}\sqrt{1-t}$. The γ_p are all positive and have the property that for any $p, q > 0$ the inequality $\sum_{i_1+\dots+i_p=q} \gamma_{i_1} \cdot \dots \cdot \gamma_{i_p} \leq \gamma_q$ holds. A power series $\sum_p a_p t^p$ with nonnegative coefficients a_p converges near zero, if and only if there exist constants $C, R > 0$ such that $a_p \leq \gamma_p \cdot C \cdot R^p$, for all $p \geq 0$.

For $n \geq 1$, let $g_n : N^{\odot n} \rightarrow M$ be the map defined as in the proof of Proposition 1.2. The only thing to verify is that the formal map $\sum_q \frac{1}{q!} \tilde{g}_q : N^{\geq 0} \rightarrow M^{\geq 0}$ strictly converges. Without restriction, we may assume that M and N are positively graded. By assumption, there are real numbers $C, R > 0$ such that $\frac{1}{p!} |f_p|^{0,\epsilon} \leq \gamma_p \cdot C \cdot R^p$, for each p . Chose $C', R' > 0$ such that

$$\|g_1\|^{0,\epsilon} \leq \gamma_1 \cdot C' \cdot R'$$

and

$$\left(\sum_{p=2}^{\infty} \gamma_p (C' \cdot R)^{p-1} \right) \cdot \|g_1\|^{0,\epsilon} \cdot C \cdot R \leq 1.$$

We show by induction on q that $\frac{1}{q!} \|\tilde{g}_q\|^{0,\epsilon} \leq \gamma_q \cdot C' \cdot (R')^q$, for each q . The case $q = 1$ is already done. For $q \geq 2$, we get:

$$\begin{aligned} \frac{1}{q!} \|\tilde{g}_q\|^{1,\epsilon} &\leq \\ \sum_{p=2}^n \sum_{i_1+\dots+i_p=n} \frac{1}{p!T!} \|g_1 \circ f_p \circ (g_{i_1} \otimes \dots \otimes g_{i_p})\|^{0,\epsilon} &\leq \\ \sum_{p=2}^n \sum_{i_1+\dots+i_p=n} \frac{1}{p!T!} \|g_1\|^{0,\epsilon} |f_p|^{0,\epsilon} \|g_{i_1}\|^{0,\epsilon} \cdot \dots \cdot \|g_{i_p}\|^{0,\epsilon} &\leq \\ \sum_{p=2}^n \sum_{i_1+\dots+i_p=n} \|g_1\|^{0,\epsilon} \gamma_p C R^p \gamma_{i_1} \cdot \dots \cdot \gamma_{i_p} (C')^p (R')^q &\leq \\ (\sum_{p=2}^n \gamma_p (C' R)^{p-1}) \|g_1\|^{0,\epsilon} C R C' \gamma_q (R')^q &\leq \\ \gamma_q C' (R')^q. \end{aligned}$$

□

4.2 Decomposition of Palamodov DG Lie algebras

In this section, we will prove our main result, an analytic version of the Decomposition Theorem 3.8. To this aim, suppose that $L = (L, d, [\cdot, \cdot])$ is a Palamodov Lie algebra, equipped with a splitting η of d such that each component $\eta^i : L^i \rightarrow L^{i-1}$ is a morphism of Palamodov spaces. In general, it is very difficult to construct such a splitting. For the case, where L is the tangent complex of a complex space, this was

realized by Palamodov. Generalisations of Palamodov's splitting construction can be found in [2]. In the Banach situation, the construction is much easier.

Again, without loss of generality, we may assume that $\eta^2 = 0$, $\eta d\eta = \eta$ and $d\eta d = d$. From now on, L stands for the quadruple $(L, d, [\cdot, \cdot], \eta)$. We define the L_∞ -structure on $H = \text{Fix}(1 - [d, \eta])$ as the L_∞ -morphism $f : H \longrightarrow L$ exactly in the same way as in Theorem 3.5 and 3.7. Both, H and the complement F of H in L inherit the graded Palamodov space structure of L . First, we show that $H = (H, \mu_*)$ is a Palamodov L_∞ -algebra.

Lemma 4.8. *Let E be a Palamodov space, $\phi \in \text{Ot}(n)$ a binary tree with n leaves and $g_1, \dots, g_{n-1} \in \text{Mult}_2(E, E)$ bilinear forms. For $\epsilon \in (0, 1)$ and each $i = 1, \dots, n-1$, suppose that $b_i := \|g_i\|^{0, \epsilon}$ and $c_i := \|\phi_i\|^{1, \epsilon}$ are finite. Then, we have the following inequalities:*

$$\begin{aligned} \|\phi(g_1, \dots, g_{n-1})\|^{0, \epsilon} &\leq b_1 \cdot \dots \cdot b_{n-1} \\ \|\phi(g_1, \dots, g_{n-1})^\sim\|^{1, \epsilon} &\leq (n-1)^{n-1} c_1 \cdot \dots \cdot c_{n-1} \end{aligned}$$

Proof. The first inequality is easy. We prove the second one by induction on n . There is only one tree β with two leaves and $\beta(g_1) = g_1$. This proves the case $n = 2$. Suppose that the statement is true, for each $m \leq n-1$. We have to show that for fixed $\lambda_0 < \lambda_1$ in $[1 - \epsilon, 1)$ and for each element $x \in E$, we have

$$(\lambda_1 - \lambda_0)^{n-1} \|\phi(g_1, \dots, g_{n-1})^\sim(x)\|_{\lambda_0} \leq (n-1)^{n-1} c_1 \cdot \dots \cdot c_{n-1} \cdot \|x\|_{\lambda_1}^n. \quad (4.15)$$

We have $\phi(g_1, \dots, g_{n-1}) = g_1 \circ \psi_1(g_2, \dots, g_p) \otimes \psi_2(g_{p+1}, \dots, g_{n-1})$, for some $p \leq n-2$ and trees $\psi_1 \in \text{Ot}(p)$ and $\psi_2 \in \text{Ot}(q)$. We will just write ϕ instead of $\phi(g_1, \dots, g_{n-1})^\sim$, ψ_1 instead of $\phi(g_2, \dots, g_k)^\sim$ and ψ_2 instead of $\phi(g_{k+1}, \dots, g_{n-1})^\sim$. Set $\lambda := \lambda_0 + \frac{1}{n-1}(\lambda_1 - \lambda_0)$. Then, we get

$$\begin{aligned} &(\lambda_1 - \lambda_0)^{n-1} \cdot \|\phi(x)\|_{\lambda_0} = \\ &(n-1)^{n-1} (\lambda - \lambda_0) \|g_1(\psi_1(x), \psi_2(x))\|_{\lambda_0} \leq \\ &c_1 (n-1)^{n-1} (\lambda - \lambda_0)^{n-2} \|\psi_1(x)\|_\lambda \|\psi_2(x)\|_\lambda = \\ &c_1 \frac{(n-1)^{n-1}}{(n-2)^{n-2}} (\lambda_1 - \lambda)^{p-1} \cdot \|\psi_1(x)\|_\lambda \cdot (\lambda_1 - \lambda)^{q-1} \cdot \|\psi_2(x)\|_\lambda \leq \\ &c_1 \frac{(n-1)^{n-1}}{(n-2)^{n-2}} \|\psi_1\|^{1, \epsilon} \cdot \|x\|_{\lambda_1}^p \cdot \|\psi_2\|^{1, \epsilon} \cdot \|x\|_{\lambda_1}^q. \end{aligned}$$

By the induction hypothesis, the last term is lower or equal

$$\begin{aligned} &c_1 \frac{(n-1)^{n-1}}{(n-2)^{n-2}} (p-1)^{p-1} c_2 \cdot \dots \cdot c_p (q-1)^{q-1} c_{p+1} \cdot \dots \cdot c_{n-1} \cdot \|x\|_{\lambda_1}^n \leq \\ &(n-1)^{n-1} c_1 \cdot \dots \cdot c_{n-1} \cdot \|x\|_{\lambda_1}^n. \end{aligned}$$

□

Theorem 4.9. *The L_∞ -algebra (H, μ_*) is Palamodov and the L_∞ -maps $f : H \longrightarrow L$ and $g : F \longrightarrow L$ are morphism of Palamodov L_∞ -algebras.*

Proof. The proofs of all three statements go almost in the same way, thus we only prove the first statement. Set $c := \|\eta\|^{0,\epsilon}$ and $k := \|[\cdot, \cdot]\|^{1,\epsilon}$. By Lemma 4.8, we have

$$\|\phi([\cdot, \cdot], \eta[\cdot, \cdot], \dots, \eta[\cdot, \cdot])\|^{1,\epsilon} \leq (n-1)^{n-1} k^{n-1} c^{n-2},$$

hence

$$\begin{aligned} \|\tilde{\mu}_n\|^{1,\epsilon} &\leq n! \left(\frac{n-1}{2}\right)^{n-1} \# \text{Ot}(n) \cdot k^{n-1} \cdot c^{n-2} \\ &\leq n!(n-1)^{n-1} \cdot (2k)^{n-1} \cdot c^{n-2} < \infty. \end{aligned}$$

The last step is by Lemma 2.2. This proves the first condition for the convergence of μ_* . To prove the second condition, set

$$\kappa := |(\text{co})\text{restriction of } [\cdot, \cdot] \text{ to } L^{\geq 1} \otimes L^{\geq 1}|^{0,\epsilon},$$

which is finite by the assumption that L is Palamodov. By Lemma 4.8 and 2.2, we get

$$\|\tilde{\mu}_n\|^{0,\epsilon} \leq \frac{n!}{2^{n-1}} \# \text{Ot}(n) \kappa^{n-1} c^{n-1} = 2^{n-1} n! \kappa^{n-1} c^{n-2}.$$

Thus, for small $t > 0$, the series $\sum_n \frac{1}{n!} \|\tilde{\mu}_n\|^{0,\epsilon} \cdot t^n$ converges. $\square$

Applying the inverse function theorem, we finally get the main result:

Corollary 4.10. *If $(L, d, [\cdot, \cdot], \eta)$ is a Palamodov DG Lie algebra with splitting η , then we have a decomposition*

$$(L, d, [\cdot, \cdot]) \cong (H, \mu_*) \oplus (F, d)$$

in the category of Palamodov L_∞ -algebras.

References

- [1] Sergey Barannikov and Maxim Kontsevich: *Frobenius manifolds and formality of Lie algebras of polyvector fields*, Int. Math. Res. Not., no. 4 (1998), 201-215.
- [2] Jürgen Bingener; Siegmund Kosarew: *Modulräume in der analytischen Geometrie*, Vieweg (1987).
- [3] M. Gugenheim; J.D. Stasheff: *On perturbations and A_∞ -structures*, Bul. Soc. Math. Belg. A38 (1987), 237-246.
- [4] Johannes Huebschmann; Jim Stasheff: *Formal solution of the master equation via HPT and deformation theory*, Forum Math. 14, no. 6 (2002), 847-868.

- [5] Tornike Kadeishvili: *On the homology theory of fibre spaces*, (Russian, English) Russ. Math. Surv. 35, no. 3, (1980), 231-238; translation from Usp. Mat. Nauk 35, no.3 (1980), 183-188.
- [6] Tornike Kadeishvili: *Algebraic structure in the homologies of an $A(\infty)$ -algebra*. (Russian. English summary) Soobshch. Akad. Nauk Gruz., SSR 108 (1982), 249-252.
- [7] Maxim Kontsevich: *Topics in algebra-deformation theory*, Berkley lectures (1995), notes by A. Weinstein.
- [8] Maxim Kontsevich: *Deformation quantization of Poisson manifolds*, Lett. Math. Phys. 66, No.3, 157-216 (2003).
- [9] Maxim Kontsevich; Yan Soibelman: *Homological mirror symmetry and torus fibrations*, Fukaya (ed.), et al., Symplectic geometry and mirror symmetry, World Scientific, (2001), 203-263.
- [10] Siegmund Kosarew: *Local moduli spaces and Kuranishi maps*, Manuscr. Math. 110, No.2 (2003), 237-249.
- [11] Lada, Tom; Markl, Martin *Strongly homotopy Lie algebras*, Commun. Algebra 23, no.6 (1995), 2147-2161.
- [12] Kenji Levêfre-Hasegawa: *Sur les A_∞ -catégories*, Thèse Université Paris 7, (2003).
- [13] Marco Manetti: *Deformation theory via differential graded Lie algebras*, Seminari di Geometria Algebrica 1998-1999, Scuola Normale Superiore (1999).
- [14] Martin Markl: *Homotopy algebras are homotopyalgebras*, Forum Math. 16, no. 1 (2004), 129-160.
- [15] S.A. Merkulov: *Operads, deformation theory and F-manifolds*, preprint AG/0210478.
- [16] S.A. Merkulov: *Strong homotopy algebras of a Kähler manifold*, Int. Math. Res. Not., no. 3 (1999), 153-164.
- [17] S.A. Merkulov: *Frobenius ∞ invariants of homotopy Gerstenhaber algebras I*, Duke Math. J. 105, (2000), 411-461.
- [18] Palamodov, V.P.: *Deformations of complex spaces*, Several complex variables. IV. Algebraic aspects of complex analysis, Encycl. Math. Sci. 10, 105-194 (1990); translation from Itogi Nauki Tekh., Ser. Sovrem. Probl. Mat., Fundam. Napravleniya 10, 123-221 (1986).

- [19] Palamodov, V.P.: *Deformations of complex spaces*, Russ. Math. Surv. 31, No.3, 129-197 (1976).
- [20] Frank Schuhmacher: *L-infini déformations et cohomologie de Hochschild*, Thèse, Institut Fourier Grenoble (2004).
- [21] Jim Stasheff: *Homotopy associativity of H-spaces*, Trans. Amer. Math. Soc., 108, (1963), 275-312.